

## § 529.40

- 529.50 Amikacin sulfate intrauterine solution.
- 529.400 Chlorhexidine tablets and suspension.
- 529.469 Competitive exclusion culture.
- 529.1003 Flurogestone acetate-impregnated vaginal sponge.
- 529.1030 Formalin.
- 529.1044 Gentamicin sulfate in certain other dosage forms.
- 529.1044a Gentamicin sulfate intrauterine solution.
- 529.1044b Gentamicin sulfate solution.
- 529.1115 Halothane.
- 529.1186 Isoflurane.
- 529.1526 Nifurpirinol capsules.
- 529.1660 Oxytetracycline.
- 529.1940 Progesterone intravaginal inserts.
- 529.2090 Salicylic acid.
- 529.2150 Sevoflurane.
- 529.2464 Ticarcillin powder.
- 529.2503 Tricaine methanesulfonate.

AUTHORITY: 21 U.S.C. 360b.

SOURCE: 40 FR 13881, Mar. 27, 1975, unless otherwise noted.

## § 529.40 Albuterol.

(a) *Specifications.* A net weight of 6.7 grams of formulated albuterol sulfate is supplied in a pressurized aluminum canister within an actuator system equipped with a detachable nasal delivery bulb.

(b) *Approvals.* See No. 000010 in § 510.600(c) of this chapter for uses as in paragraph (d) of this section.

(c) *Special considerations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d) *Conditions of use—(1) Amount.* Each valve actuation (puff) of the device delivers 120 micrograms (mcg) of albuterol sulfate. One dose is three (3) puffs, totaling 360 mcg.

(2) *Indications for use.* For the immediate relief of bronchospasm and bronchoconstriction associated with reversible airway obstruction in horses.

(3) *Limitations.* Not for use in horses intended for food.

[67 FR 7072, Feb. 15, 2002]

## § 529.50 Amikacin sulfate intrauterine solution.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 250 milligrams of amikacin (as the sulfate).

(b) *Sponsor.* See No. 000856 and 059130 in § 510.600(c) of this chapter.

## 21 CFR Ch. I (4–1–05 Edition)

(c) *Conditions of use—(1) Amount.* Two grams (8 milliliters) diluted with 200 milliliters of sterile physiological saline per day for 3 consecutive days.

(2) *Indications for use.* For treating genital tract infections (endometritis, metritis, and pyometra) in mares when caused by susceptible organisms including *E. coli*, *Pseudomonas* spp., and *Klebsiella* spp.

(3) *Limitations.* For intrauterine infusion in the horse only. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 9640, Mar. 8, 1983, as amended at 53 FR 27852, July 25, 1988; 62 FR 15110, Mar. 31, 1997; 62 FR 23358, Apr. 30, 1997]

## § 529.400 Chlorhexidine tablets and suspension.

(a) *Specification.* Each tablet and each 28-milliliter syringe of suspension contain 1 gram of chlorhexidine dihydrochloride.<sup>1</sup>

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* Place 1 or 2 tablets deep in each uterine horn; or infuse a solution of 1 tablet dissolved in an appropriate amount of clean boiled water; or infuse one syringe of suspension into the uterus.<sup>1</sup>

(2) *Indications for use.* For prevention or treatment of metritis and vaginitis in cows and mares when caused by pathogens sensitive to chlorhexidine dihydrochloride.<sup>1</sup>

(3) *Limitations.* Prior to administration, remove any unattached placental membranes, any excess uterine fluid or debris, and carefully clean external genitalia. Use a clean, sterile inseminating pipette for administering solutions and suspensions. Treatment may be repeated in 48 to 72 hours.<sup>1</sup>

[43 FR 10705, Feb. 23, 1979]

## § 529.469 Competitive exclusion culture.

(a) *Specifications.* Each packet of lyophilized culture contains either

<sup>1</sup>These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.